

# ATTENTION DRUG DISPOSAL STAFF

## Rx Destroyer™ Guidelines:

1. Document waste in your Pyxis, Omnicell, controlled drug inventory sheet, or other automated medication dispensing cabinet.
2. Load medications into the bottle CII-CV-Funnel use is optional.
  - Load pills, suppositories, powders, creams, IV BAG waste.
  - Fold patches before inserting into bottle.
  - Squirt syringe dose into bottle. -- DO NOT place syringe or vial into the bottle, use sharps container.
3. Tightly secure cap and gently agitate for 10 seconds. (recommended at least 1-2 times/day or per usage)
4. After each drug disposal session, repeat step #2.
5. Bottle is full when contents rise within 2" from top of bottle.
6. NOTIFY your internal designated department to remove the bottle to be placed in the proper disposal bin, according to internal discard process.
7. NOTIFY your internal designated department to replace the new Rx Destroyer™ bottle securely into the Rx Destroyer™ locking accessory.



**#1** in Hospitals  
& Surgery  
Centers

### Questions/Contact Info:

Rx Destroyer™ 888-608-5160 • [www.rxdestroyer.com](http://www.rxdestroyer.com)

### Warnings:

#### No effervescent medications

Refer to state and local hazardous waste disposal regulation • DO NOT INGEST will induce vomiting • If ingested drink large amounts of water or milk: CONSULT A PHYSICIAN IMMEDIATELY • First aid for eyes, flush with cool water for 15 minutes: CONSULT A PHYSICIAN IMMEDIATELY • Keep out of reach of children • Bottle contents may freeze – Store above 40 degrees

**DISCLAIMER:** Information for reference only. Users of Rx Destroyer™ products agree to perform due diligence in understanding and abide by specific process, rules and laws by which they are governed. The generator accepts sole responsibility for proper inserted contents and witness documentation of contents inside the Rx Destroyer™